

The underscreened Kondo effect in ladder systems

Karyn Le Hur

Theoretische Physik, ETH-Hönggerberg, CH-8093 Zürich, Switzerland

We introduce an unfamiliar Kondo ladder problem by coupling the well-known Takhtajan-Babujan $S=1$ chain to a half-filled one-dimensional electron gas, and we solve it using quantum field theory techniques. Based on the so-called underscreened Kondo effect, we find an insulating spin liquid with both deconfined spinons and optical magnons. Important consequences are discussed.

PACS numbers: 71.10.Pm, 71.27.+a, 75.10.Jm

The problem of the co-existence between conduction electrons and an array of local magnetic moments remains an eminent unsolved problem in condensed matter physics, in link with the well-known Kondo effect. This poses a real challenge to the theorist. As a first step, the one-dimensional (1D) model of a Kondo chain with an M -fold degenerate band of conduction electrons interacting with a spin S Heisenberg chain has been solved in a semi-classical limit [1]. At low energy, the system is insulating and the spin properties can be described by an $O(3)$ nonlinear sigma model ($NL\sigma M$) with a topological term $\theta = \pi(2S - M)$. For $|M - 2S| = (\text{even})$, single electron excitations at low energies are massive spin polarons. For $|M - 2S| = (\text{odd})$, the $NL\sigma M$ in contrast becomes critical and belongs to the universality class of the isotropic $S=1/2$ Heisenberg chain. In this Letter, we investigate the interesting case $S=1$ and $M=1$ in a so-called ‘ladder’ picture [2]. As we shall see below, the characterization of the spin array as an $S=1$ quantum spin chain with prevalent gapless triplet excitations should allow to push forward our understanding of the underscreened Kondo effect [3] in the Kondo lattice model (KLM). For experimental applicability, this model appears to be particularly appropriate to describe heavy-fermion materials with Uranium compounds such as UPt_3 (as pointed out in ref. [4], where we have firstly solved the problem of two localized $S=1$ spins embedded in a host metal).

Let us start with the integrable Takhtajan-Babujan $S=1$ chain on a lattice, described by: $\mathcal{H} = J_{\parallel} \sum_j (\mathbf{S}_j \mathbf{S}_{j+1}) - \beta (\mathbf{S}_j \mathbf{S}_{j+1})^2$ where $\beta = 1$ [5]. $\mathbf{S}_j$ describe local moments. The unusual term quartic in spin can be generated, for example, by phonons [6]. Solving Bethe Ansatz equations, the only elementary excitation is known to be a doublet of gapless spin-1/2 spin waves, with total spin 0 or 1 [5]. The approach discussed in this Letter is based on field theory techniques like the non-Abelian [7] and Abelian [8] bosonizations. Thus, it is convenient to use a (simple) *continuum* description of such an integrable model. Interestingly, the specific heat behaves as: $C_v/L = 2ST/(1 + S) + \mathcal{O}(T^3)$, where $S=1$ [5]. In the sense of critical theories, such a model could be parametrized by a conformal anomaly (which is the central charge in the Virasoro algebra): $C = \frac{3S}{S+1} = 3/2$ [9]. This fact, together with approximate

mappings, leads to the conjecture that the criticality of this model is governed by operators satisfying a critical Wess-Zumino-Witten (WZW) model on the $SU(2)$ group at the level $k = 2S = 2$ [10]. Solving numerically the Bethe-Ansatz equations the authors of ref. [11] checked the main conjectures made in refs. [9,10]. Thus, for $\beta = 1$, the effective action for the low-energy excitations is given by the WZW model on the $SU(2)$ group at the level $k = 2$. The following general statement is valid: the Kac-Moody algebra for the group $SU(N)$ at the level $k=N$ can be represented by the currents of Majorana fermions $\xi^a (a = 1, \dots, N^2 - 1)$ from the adjoint representation of this group. Thus, the $SU(2)_{k=2}$ currents can be explicitly rewritten as: $I^a(\bar{I}^a) = -\frac{i}{2} \epsilon^{abc} \xi_{L(R)}^b \xi_{L(R)}^c$. The Hamiltonian is in turn equivalent to the model of three massless Majorana fermions $\xi^a (a = 1, 2, 3)$ [12]. In the continuum limit, the spin-1 chain has *only* zero sound triplet excitations (or *magnons*, like in conventional magnets). The staggered magnetization $\mathbf{N}$ can be simply expressed as $\mathbf{N}(x) \simeq \text{Tr}\{(h + h^\dagger)\sigma\}$, where $h \in SU(2)_{k=2}$ has dimension $3/8$ [10,12].

The spectrum of *free* conduction electrons yields a separation of spin and charge. The spin sector is described by an $SU(2)$ WZW model with $k=1$ [7]. Here, the two fundamental fields $g, g^\dagger$ are *spinon* pairs which can be viewed as free fields except from a purely statistical (in that case: semionic) interaction. The electronic spin density is usually described as: $\mathbf{S}_c(x) = \mathbf{J}_c(x) + e^{2ik_F x} \mathbf{N}_c(x)$, where $\mathbf{J}_c = \mathbf{J} + \bar{\mathbf{J}}$ and $\mathbf{N}_c \simeq \text{Tr}\{\sigma(g + g^\dagger)\} \cos \sqrt{2\pi} \Phi_c^e$ are, respectively, the smooth and staggered parts of the magnetization. $\mathbf{N}_c$ has the dimension 1 and $\mathbf{J}_c$ can be explicitly written as a function of g and $g^\dagger$ [7]. The charge sector is described in terms of a $U(1)$ scalar field Φ_c^e leading to a Conformal Field Theory (CFT) with $C=1$ for holons as well. Zero sound charge modes propagate with the velocity v_ρ and are ruled by the Luttinger parameter $K_\rho = v_F/v_\rho$. For a 1D free electron gas $K_\rho = 1$, and $v_F = 2t \sin(k_F a)$ is the velocity for charge and spin degrees of freedom. The Kondo interaction takes the form:

$$\mathcal{H}_{int} = \lambda_2 (\mathbf{J} + \bar{\mathbf{J}})(\mathbf{I} + \bar{\mathbf{I}}) + \alpha \lambda_3 e^{i\delta x} \text{Tr}(g^\dagger \sigma) \cos \sqrt{2\pi} \Phi_c^e \mathbf{N} + h.c. \quad (1)$$

where $\lambda_{2,3} \propto J_K$, and α is a non-universal constant.

We have defined $\delta = 2k_F - \pi$. Here, we consider situations of the half-filled case, with $\delta \rightarrow 0$. The bare conditions are fixed to: $J_K \ll (J_{||}, t)$. The coupling between the short-range staggering fluctuations becomes there prominent. A small term λ_3 modifies the universal dc conductivity of the electron gas $\sigma_o = 2e^2 K_\rho L/h$, obtained by applying a static field over a finite part of the pure sample. In the presence of magnetic impurities, the dc conductivity is rescaled as $\sigma(T) \propto \{T\lambda_3(T)^2\}^{-1}$ [13]. We assume that the phase coherence is lost after each magnetic collision. Then, the diffusion by the array of magnetic impurities with S=1 provides an anomalous power-law, $\sigma(T) \propto T^{1/4}$. Note that the smallest amount of Kondo ‘disorder’ will destroy the so-called Umklapp term which gives $\sigma(T) \sim T^{-1}$ (when $K_\rho = 1$) by using a memory-function approximation [14]. Clearly, the effects of scattering from the array of magnetic impurities become coherent at lower energy which typically give rise to a Kondo localization. An estimate for the localization length may be simply obtained from one-loop perturbation theory; it gives: $\mathcal{L}_{mag} = (1/J_K)^{8/5}$.

The dc conductivity for this state can be easily computed using the Kubo formula. The resistivity follows $\rho(T) \propto e^{\Delta/k_B T}$ when $k_B T \ll \Delta$ leading to a perfect insulating state. When $\delta = 0$, the mass gap (which can be identified either as the charge gap or the spin gap since λ_3 couples spin and charge sectors) obeys $\Delta = \mathcal{L}_{mag}^{-1} \equiv J_K^{8/5}$. It is larger than in the usual 1D KLM (with S=1/2) for the same bare conditions [13]. The only possibility to obtain a non-zero conductivity is through thermal fluctuations which can activate charge excitations. When $T \rightarrow 0$, correlations of $\cos \sqrt{2\pi} \Phi_c^c$ show a long-range order. Summing the spin part and refermionizing the theory, the diagonalization of the charge sector is an easy task even for $\delta \neq 0$. We find a gas of spinless fermions ruled by the dispersion relation: $\epsilon(k) = \pm \sqrt{v_F^2 k^2 + \Delta^2} - v_F \delta/2$. It rescales the mass gap as $\Delta(\delta) = \Delta - v_F \delta/2$ and the insulating state can be stabilized as long as $\Delta \gg v_F \delta/2$.

Now, let us investigate the spin properties of such an insulating Kondo system (for $\delta = 0$) and explain the crucial idea of the paper. The physics behind these calculations is the formation of bound states between conduction electrons and localized spins, leading to a semi-conducting phase. Of course, the occurrence of Kondo bound states can be achieved only due to the splitting of each S=1 localized spin onto two doublet modes. Note that although powerful techniques have been used to solve the S=1 impurity Kondo model, it is difficult to have a ‘simple’ mathematical formulation of the splitting phenomenon (see ref. [4] and references therein). In the present lattice model, we express it simply as follows. For $T \simeq \Delta$ (typically the spin gap) we use the decomposition,

$$\mathbf{N} \Delta^{1/8} = (\mathbf{N}_1 + \mathbf{N}_2) \quad (2)$$

with $\mathbf{N}_i = \text{Tr}(h_i \sigma)$ and $h_i \in SU(2)_{k=1}$. Through the fac-

tor $\Delta^{1/8}$, we consider that the operators h_i (with $k=1$) have the scaling dimension 1/2 and not 3/8. Only ‘half’ of each S=1 spin can form a bound state with a conduction electron. Then, one doublet is assumed to participate in the Kondo process whereas the other would remain massless. This phenomenon can be simply described through the important formula:

$$\frac{\mathbf{N}}{2} \text{Tr}(g\sigma) = \Delta^{-1/8} \text{Tr}(h_1^\dagger \sigma) \text{Tr}(g\sigma) \quad (3)$$

First, the SU(2) field h_2 remains free (except statistical interactions), leading to a critical C=1 CFT (i.e. a Heisenberg model) with deconfined spinons in the ground state. Then, despite of the insulating state the optical conductivity would reveal only a pseudogap at $k = \pi$ through spin instantons. Following the arguments of Tsvelik [1], one can define an anomalous fermionic current $\mathbf{J}_A$ which is proportional to the topological charge density of the staggered magnetization $\mathbf{N}_2$. Thus, $\mathbf{J}_A$ varies linearly with $\frac{1}{\pi} \epsilon^{\mu\nu} \mathbf{N}_2 (\partial_\mu \mathbf{N}_2 \times \partial_\nu \mathbf{N}_2)$ which has the scaling dimension 5/2. In the (q, w) space, it gives a purely dynamical optical conductivity $\sigma_{opt}(w, k = \pi - q) \propto \frac{1}{w} (w^2 - v_F^2 q^2)^{3/2} \sim w^2$. Second, deconfinement of spinons does not imply that conventional magnons fail to be quasiparticles in such a spin liquid. The low-energy physics also involves optical magnons. It is convenient to define: $\eta = \lambda_3 \langle \cos \sqrt{2\pi} \Phi_c^c(x) \rangle \propto \Delta^{9/8}$. During the spin-1 splitting mechanism, the Kondo exchange is rescaled; we change, $2\eta \rightarrow \eta$. Then, by using Eq.(3), the dimensional equality $\eta_0 = \eta \Delta^{-1/8} \simeq \Delta$ and omitting high-energy singlet magnon modes (see later), we find a massive part described by:

$$S_U = \int d^2x \frac{1}{2\gamma} \text{Tr}(\nabla U \nabla U^\dagger) - \eta_0 : \text{Tr}(U + U^\dagger) : \quad (4)$$

where, $\gamma^{-1} \propto \sqrt{v_F/J_{||}a} + \sqrt{J_{||}a/v_F}$, $x_0 = \tau \sqrt{v_F J_{||}a}$, $x_1 = x$ and $U(x) = gh_1^\dagger(x)$ has SU(2) symmetry.

Excitations of the O(3) NL σ M are S=1 triplets (magnons). At low temperatures, $\text{Tr}U(x) = \text{constant}$ and then magnons become purely massive. Since by construction $\text{Tr}U$ has the scaling dimension 1, the spin gap (of course) is of order of $\eta_0 \simeq \Delta$. The action S_U can be rewritten as a triplet of Majorana fields $\chi^a (a = 1, 2, 3)$ with the same mass $m_t = \Delta$. They mainly contribute to a coherent δ -peak to the dynamical susceptibility $\chi''(q = \pi - k > 0, \omega) \sim m_t (q^2 + m_t^2)^{-1/2} \delta(\omega - \sqrt{q^2 + m_t^2})$ near $k = \pi$ and $\omega = m_t$. On the other hand, the spectral density of the staggered magnetization $(-1)^x \mathbf{N}_2(x)$ with $\mathbf{N}_2 \sim \text{Tr}(h_2 \sigma)$ shows a purely incoherent background. Schulz [15] explained that it leads to: $\chi''(q > 0, \omega) \simeq \sin(2\pi c) T^{4c-2} \Im m[\rho(\frac{w-q}{4\pi T}) \rho(\frac{w+q}{4\pi T})]$ with $\rho(x) = \frac{\Gamma(c-ix)}{\Gamma(1-c-ix)}$ and $c=1/4$ for the isotropic Heisenberg chain. Such a contribution diverges when $\omega \rightarrow q$ as $(w - q)^{-1}$ so that the δ -peak due to optical magnons remains distinguishable at $w = \sqrt{q^2 + m_t^2}$.

During the Kondo localization, the current $\mathbf{I}$ is also decomposed into two independent parts $(\mathbf{J}_1, \mathbf{J}_2)$ where $\mathbf{J}_i$ are $SU(2)$ currents at the level $k=1$ (with $C=1$ each). Some degrees of freedom should account for the missing $C=1/2$. Those are associated with the discrete (Z_2) nonmagnetic interchange symmetry $1 \leftrightarrow 2$ in the definition of $\mathbf{I}$. The marginal term can be rewritten: $\lambda_2(\mathbf{J}_c + \bar{\mathbf{J}}_c)(\mathbf{J}_1 + \bar{\mathbf{J}}_1)$. We have the correspondence $(\mathbf{J}_1 \oplus \mathbf{J}_c) = -\frac{i}{2}\epsilon^{abc}\chi_L^b\chi_L^c$. Then, by using the Dyson equation for the fields χ_i , we find that the mass m_t is slightly renormalized: $\Delta(1 + \frac{J_K a}{2\pi v_F} \ln \frac{\Lambda}{\Delta})$ and $\Lambda \sim t$. Next, we neglect these logarithmic corrections. For $J_{\parallel}a \neq v_F$, free magnon excitations can survive for $T < \Delta$. We can integrate out excitations which correspond to configurations where $\text{Tr}U \neq 0$ acquires a gap. It becomes an $O(3)$ NL σ M where the topological term has no contribution. Finally, it provides a rescaled mass gap $m_t \sim \Delta \exp(-2\pi\tilde{\gamma}^{-1})$, where $\tilde{\gamma}^{-1} \propto |J_{\parallel} - t|$. Here, the spin correlation functions of the massive part are proportional to the Mac-Donald function $K_0(m_t \mathbf{r})$ with $\mathbf{r} = (x_0, x_1)$ and $K_0(x) = \sqrt{\frac{\pi}{2}}x^{-1/2}e^{-x}(1 - 1/8x + \mathcal{O}(x^{-2}))$ [16]. Massive polarons should occur in the interval between m_t and Δ due to interactions of electrons and kinks in $\mathbf{N}_1$ [1].

The conclusion that the fixed point is the $k=1$ WZW model is consistent with Zamalodchikov's c-theorem [17]. This theorem states that if a field theory flows between two conformally invariant fixed points, then the value of the conformal anomaly parameter C must be smaller at the Infra Red stable fixed point. Here, the unstable fixed point is ruled by $C = 7/2$; the stable fixed point has $C = 1$. Deconfined spinons remain as a result of the non-Abelian chiral symmetry which is not broken. Note that the spinon velocity $v \sim J_{\parallel}a$ is equal to that in the single $S=1/2$ spin chain problem [18]. Through the transform $\xi^i \rightarrow \chi^i$, magnons become optical. For $T \ll \Delta$, only the gapless modes (spinons) contribute to the specific heat $C_v = \pi T/3v$. Now, we choose the particular point: $J_{\parallel}a = v_F$. Using the Abelian bosonization we proceed to calculate quantities of interest like the spin-spin correlations, the NMR relaxation rate $\mathcal{T}_1$ and the magnetization by varying the external magnetic field.

First, notice that an alternative picture of the integrable $S=1$ chain with $\beta = 1$ is realized in the context of spin ladders, when adding a strong four-spin interaction $V \sum_j \mathbf{S}_1(j)\mathbf{S}_1(j+1)\mathbf{S}_2(j)\mathbf{S}_2(j+1)$ equal to the direct transverse exchange $J_{\perp} < 0$ [19]. Second, using the common bosonization scheme for the two-leg ladder (see, for instance, Parts II and III of ref. [20]), we can rewrite the magnetization $\mathbf{N}$ of the $S=1$ chain in terms of two scalar fields Φ_e and Φ_o , and their duals Θ_e and Θ_o , as follows:

$$\begin{aligned} N_x &= \cos \sqrt{\pi}\Theta_e \cos \sqrt{\pi}\Theta_o \\ N_y &= \sin \sqrt{\pi}\Theta_e \cos \sqrt{\pi}\Theta_o \\ N_z &= \sin \sqrt{\pi}\Phi_e \cos \sqrt{\pi}\Phi_o \end{aligned} \quad (5)$$

For $J_{\perp} = V$, note that Φ_e and Θ_e are completely free,

whereas the Hamiltonian of Φ_o describes a single free Majorana field. It leads to a CFT with a total central charge $C=3/2$. The operators $\exp i\sqrt{\pi}\Phi_e(\Theta_e)$ and $\cos \sqrt{\pi}\Phi_o(\Theta_o)$ acquire the scaling dimensions $1/4$ and $1/8$, respectively. Now, we use the representation

$$\mathbf{N}_c = (\cos \sqrt{2\pi}\Theta_c, \sin \sqrt{2\pi}\Theta_c, \sin \sqrt{2\pi}\Phi_c) \quad (6)$$

Then, it is not difficult to show that the splitting of each $S=1$ local spin can be still understood through the formula (2) where $\mathbf{N}_{i=1,2}$ has the same form as $\mathbf{N}_c$, but replacing $\Phi_c \rightarrow \Phi_i$ and $\Theta_c \rightarrow \Theta_i$. We have the constraints, $\Phi_{1,2} = 1/\sqrt{2}(\Phi_e \pm \Phi_o)$. By using the Abelian bosonization, Eq.(3) can be understood as follows: only one bosonic field, namely Φ_1 , can hybridize with Φ_c and similarly for the dual fields. The term λ_3 can be simply written as:

$$\begin{aligned} \mathcal{H}_{int} &= -\eta_1 : \cos \sqrt{4\pi}\Phi_+(x) : + \eta_2 : \cos \sqrt{4\pi}\Phi_-(x) : \\ &+ 2\eta_3 : \cos \sqrt{4\pi}\Theta_-(x) : \end{aligned} \quad (7)$$

with the notations: $\eta_i \propto \Delta/\pi^2 a$ and $\Phi_{+,-} = 1/\sqrt{2}(\Phi_1 \pm \Phi_c)$. The physics depicted by $\mathcal{H}_{int}$ becomes similar to that of the two-leg ladder system, with $V = 0$.

Results obtained in ref. [20] are then applicable to that case and we present them briefly for completeness. The Abelian bosonization method gives an *exact* massive spectrum which is composed of a triplet of Majorana fields $\chi^a (a = 1, 2, 3)$ with the same mass $m_t = \Delta$, and a singlet χ^o branch shifted to the high energy $m_s = -3\Delta$. The total $\mathbf{N}^{(+)}$ and relative $\mathbf{N}^{(-)}$ staggered magnetization of the *massive* part are given by,

$$\begin{aligned} \mathbf{N}^{(+)} &= \{\sigma_1\mu_2\sigma_3\sigma, \mu_1\sigma_2\sigma_3\sigma, \sigma_1\sigma_2\mu_3\sigma\} \\ \mathbf{N}^{(-)} &= \{\mu_1\sigma_2\mu_3\mu, \sigma_1\mu_2\mu_3\mu, \mu_1\mu_2\sigma_3\mu\} \end{aligned} \quad (8)$$

where σ_i, μ_i are order and disorder parameters of three Ising models corresponding to the massive triplet of the Majorana fields (χ^a) while σ, μ refer to the singlet branch (χ^o). Masses determine deviations from criticality such as $m_{t(s)} \equiv (T - T_c)/T_c > (<) 0$. It corresponds to three disordered phases with $\langle \mu_j \rangle \neq 0$ and $\sigma_j = 0$, and one ordered phase with $\langle \mu \rangle \neq 0$ and $\sigma = 0$. We used the so-called Kramers-Wannier duality $t \rightarrow -t, \sigma \leftrightarrow \mu$ for (σ, μ) corresponding to a negative mass. Using the asymptotic behaviors of the different 2-point Ising correlation functions [20] and the expression of $\mathbf{N}_2$, one obtains:

$$\begin{aligned} \langle \mathbf{N}^{(-)}(\mathbf{r})\mathbf{N}^{(-)}(\mathbf{0}) \rangle &\propto \frac{1}{r^{5/2}}e^{-m_t r} + \mathcal{O}(e^{-3m_t r}) \\ \langle \mathbf{N}^{(+)}(\mathbf{r})\mathbf{N}^{(+)}(\mathbf{0}) \rangle &\simeq \frac{1}{r^{3/2}}e^{-2(m_t + m_s)r} \\ \langle \mathbf{N}_2(\mathbf{r})\mathbf{N}_2(\mathbf{0}) \rangle &\simeq \frac{1}{r} \end{aligned} \quad (9)$$

For $J_K > 0$, the $S=1$ magnetization is essentially determined by the *relative* staggered part $\mathbf{N}^{(-)} = \mathbf{N}_1 - \mathbf{N}_c$.

The singlet branch gives only high-energy corrections so that the non-Abelian action S_U becomes exact at low temperatures. The fields Φ_2 and Θ_2 remain free. Thus, N_2 yields the typical correlation functions of the $S=1/2$ isotropic Heisenberg chain. A related quantity of direct experimental relevance is the NMR relaxation rate T_1 . It is easy to see that the staggered susceptibilities dominate in T_1^{-1} . Precisely, we obtain: $\chi(2k_F) = 1/T + T^{-1/2} \exp -\Delta/T$. For a system of non-interacting spins $\chi(2k_F) = (T_1 T)^{-1}$ is a constant. Here, the two-component underlying magnetic order changes the effective field seen by the nuclei and, for instance, T_1^{-1} goes to a nonzero value as $T \rightarrow 0$.

Pleasantly, as in three-coupled spin chain systems [21], there is a phenomenon which is strikingly analogous to the quantum Hall effect. The infinite degeneracy of a Heisenberg chain is lifted for a strong magnetic field. The system shows a plateau in the magnetization curve at magnetization *per site* $m=1/2$ and remains insulating even when a finite magnetization is induced by the external field. Thus, we obtain the same physics as for a partially magnetized $S_{tot} = 3/2$ chain [22]. Regarding each $S_{tot} = 3/2$ operator as a symmetrized product of three $S=1/2$'s, one $S=1/2$ is polarized by the applied field at each site while the other two form a valence-bond-state having a nonlocal topological order associated with the breakdown of a hidden discrete $\mathcal{Z}_2 \times \mathcal{Z}_2$ symmetry. Here, we have a two-chain realization of such a Haldane state and the 'string' order parameter is defined as $\lim_{|x-y| \rightarrow +\infty} \langle \mathcal{O}_{str}(x, y) \rangle \simeq \langle \mu_1 \rangle^2 \langle \mu_2 \rangle^2 \neq 0$. But, we expect the vanishing of the Haldane gap for some value $H = H_{c1}$ of the magnetic field. We closely follow the notations of ref. [23]. We can see easily that H affects only the symmetric field Φ_+ as, $\mathcal{H}_+ \rightarrow \mathcal{H}_+ + \sqrt{2}H\partial_x\Phi_+$. As long as $H < H_{c1}$ the result is a Sine-Gordon model with a gap which is rescaled by the magnetic field. It can be viewed as a completely filled lower band for spinless fermions with a gap $\sim \Delta$ separating the lower and the upper bands. The Fermi level, lying in the middle, is shifted when increasing H . H_{c1} refers to the situation where the Fermi energy crosses the gap and just lies at the bottom of the upper band. We have the relation $\Delta = g\mu_B H_{c1}$. Above H_{c1} , the upper band is partially filled leading to an amount in the magnetization $\delta m = m - 1/2$. The problem is very similar to the commensurate-incommensurate transition [24]. Near the plateau $\delta m \propto \sqrt{|H - H_{c1}|}$ and $1/T_1 \sim T^{-1/2}$. The system remains *insulating* since Θ_- has a nonzero expectation value and Φ_- has a gap [23].

In conclusion, we have presented a new Kondo ladder system by coupling the so-called Takhtajan-Babujan $S=1$ chain to a half-filled 1D electron gas and obtained low-energy properties using quantum field theory techniques. Through the underscreened Kondo effect, this gives novel insights into 1D spin liquids: the optical conductivity reveals only a pseudogap at $k = \pi$ despite of the insulating state, a coherent magnon peak in the dynamical spin

susceptibility coexists with an incoherent spinon background, the magnetization yields a fractional-like quantization. Such a Kondo state still occurs in the vicinity of the $\beta = 1$ point, but when $J_{||}(1 - \beta) = m_t$ the $S=1$ chain is in the Haldane phase [25] and the electron gas is decoupled. These results extend the semi-classical picture of ref. [1] which is applicable only for $J_{||}a/v_F \rightarrow 0$ and should explain the coexistence between a Kondo singlet state and strong antiferromagnetic fluctuations in some heavy fermion compounds such as UPt_3 . More generally, this problem belongs to the class of Luttinger liquids in active environments, a topic of importance not only to the Kondo chain physics, but also to e.g. striped phases and carbon nanotubes. K.L.H. thanks B. Coqblin, A. Honecker, T.M. Rice and A. Tsvelik for conversations.

-
- [1] A.M. Tsvelik, Phys. Rev. Lett. **72**, 1048 (1994) and cond-mat/9405024.
 - [2] E. Dagotto and T.M. Rice, Science **271**, 618 (1996).
 - [3] P. Nozières and A. Blandin, J. Physique. (Paris) **41**, 193 (1980).
 - [4] K. Le Hur and B. Coqblin, Phys. Rev. B **56**, 668 (1997).
 - [5] L.A. Takhtajan, Phys. Lett. A **87**, 479 (1982); M. Babujan, Nucl. Phys. B **215**, 317 (1983).
 - [6] Concerning the experimental role of the biquadratic term, see: E.L. Nagaev, Usp. Fiz. Nauk. **25**, 61 (1982).
 - [7] I. Affleck, Nucl. Phys. B **265**, 409 (1986).
 - [8] F.D.M. Haldane, J. Phys. C **14**, 2585 (1981).
 - [9] I. Affleck, Phys. Rev. Lett. **56**, 764, 2763 (1986).
 - [10] I. Affleck and F.D.M. Haldane, Phys. Rev. B **36**, 5291 (1987).
 - [11] F.C. Alcaraz and M.J. Martins, Phys. Rev. Lett. **61**, 1529 (1988). The results are in agreement with the generalized Coulomb-gas picture of P. Di Francesco, H. Saleur and J.B. Zuber, Nucl. Phys. B **300**, 393-432 (1988).
 - [12] A.M. Tsvelik, Phys. Rev. B **42**, 10499 (1990).
 - [13] K. Le Hur, Phys. Rev. B **58**, 10261 (1998).
 - [14] T. Giamarchi, Phys. Rev. B **44**, 2905 (1991).
 - [15] H.J. Schulz and C. Bourbonnais, Phys. Rev. B **27**, 5856 (1983); H.J. Schulz, *ibid.*, **34**, 6372 (1986).
 - [16] A.N. Kirillov and F.A. Smirnov, Int. J. Mod. Phys. **3**, 731 (1988).
 - [17] A.B. Zamolodchikov, J.E.T.P. Lett. **43**, 730 (1986).
 - [18] This result is in complete agreement with the problem of three leg (spin) ladders: B. Frischmuth, B. Ammon and M. Troyer, Phys. Rev. B **54**, 3714 (1996).
 - [19] A.A. Nersesyan and A.M. Tsvelik, Phys. Rev. Lett. **20**, 3939 (1997).
 - [20] D.G. Shelton, A.A. Nersesyan and A.M. Tsvelik, Phys. Rev. B **53**, 8521 (1996).
 - [21] M. Oshikawa, M. Yamanaka and I. Affleck, Phys. Rev. Lett. **78**, 1984 (1997); D.C. Cabra, A. Honecker and P. Pujol, *ibid.*, **79**, 5126 (1997).
 - [22] M. Oshikawa, J. Phys. Condens. Matter **4**, 7469 (1992).

- [23] R. Chitra and T. Giamarchi, Phys. Rev. B **55**, 5816 (1997).
- [24] G.I. Dzhasharidze, A.A. Nersisyan, JETP Lett. **27**, 334 (1978); V.L. Prokovsky, A.L. Talapov, Phys. Rev. Lett. **42**, 65 (1979); H.J. Schulz, Phys. Rev. B **22**, 5274 (1980).
- [25] F.D.M. Haldane, Phys. Lett. **93A**, 464 (1983).